

CASE REPORT

Odontoma- A Benign Mixed Odontogenic Tumor –Report of Two Cases

Bharanippriya.V¹, Preethe.M², Umamaheswari.G³, Vidyalakshmi.S⁴

ABSTRACT : The term Odontoma was coined by Paul Broca in 1869, who defined it as a tumor formed by overgrowth of complete dental tissue. Odontomas constitute about 22% of the odontogenic tumours of the jaws. Here we report 2 cases in the age range of 9-12years visited our OPD with the chief complaint of missing teeth. Routine clinical, radiographic and histopathological examination revealed the presence of compound odontoma which was removed surgically. Being a benign odontogenic neoplasm early diagnosis of odontomas ensure better prognosis

Keywords: *Odontoma, Compound odontoma, Mixed odontogenic tumor, Hamartomatous lesion*

Introduction :

Odontoma is an odontogenic mixed benign neoplasm that originates from epithelial and ectomesenchymal component. They are generally small but occasionally grow to large sizes.^[1] Most authors accept that odontoma represents a hamartomatous malformation rather than a true neoplasm.^[2] Odontoma is called as composite odontoma, since it is composed of more than one type of tissue such as enamel, dentin, variable amounts of cementum and pulpal tissue.^[3,4] Aetiology of odontoma is unknown. Local trauma/infection can cause odontoma. Males show a slight higher predilection when compared to females.^[2,6] The mean age of incidence is around 14 years and most prevalent age of diagnosis is around 2nd decade of life.^[5] Maxilla shows greater predilection than mandible at the ratio of 3:1.^[2] In general odontomas are asymptomatic, occasionally presents with signs and symptoms related to their surrounding structures, such as unerupted / impacted teeth ,retained deciduous teeth ,swelling and evidence of infection.

Odontoma can be classified into three variants clinically: Central (intraosseous) odontoma, peripheral (extraosseous) odontoma and erupted odontoma. Intraosseous odontoma is further sub-classified into compound and complex odontoma. The compound type is more prevalent in maxillary anteriors (61%) and complex in (59%).^[2] Compound odontoma characteristically comprises multiple, small tooth like structures whereas complex odontoma presents as a conglomerate mass of dentin and enamel.^[5]

Case Report - 1

A 9 year old female came with the complaint of missing upper right second tooth. On intra oral examination a diffuse swelling was seen in the region of 12 which was firm to hard in consistency. Intra oral periapical radiograph in 12 revealed a radiopaque mass which had obstructed the eruption of 12 [**Figure 1a**]. Macroscopic examination revealed a small primary tooth like structure which was hard in consistency [**Figure 1b**]. Microscopic examination of the ground section of the specimen revealed well organized enamel and dentin like structures with no evidence of root [**Figure 1c**].

Case Report - 2

A 11 year old male came with the complaint of missing anterior tooth. On intraoral examination a firm swelling was noticed in maxillary anterior region in the region of 11, but it was not associated with any pain. Upon radiographic investigation, OPG revealed a radiopaque mass obstructing the eruption pathway of the impacted 11 [**Figure 2a**]. The mass is removed and patient referred to orthodontic alignment. Macroscopy revealed hard tissue bits resembling teeth with crown and root like portions. Small tooth like structure fused to form large agglomerate masses [**Figure 2b**]. Macroscopic examination of the ground section revealed a thin hypoplastic enamel and dentin like areas, a completely formed root with cementum like structure. Fusion of three teeth like structures were evident [**Figure 2c**].

Discussion :

The world health organization (WHO) defined the odontoma as ‘a malformation in which all the dental tissues are represented, individual tissues being mainly formed but occurring in more or less disorderly pattern’. It is the most commonly occurring benign odontogenic tumour (32-40%) and was coined in the year 1869 by Paul Broca.^[2] Odontoma is considered to be a mixed tumour having both epithelial and ectomesenchymal component. Now, it is considered as a developmental anomaly or hamartomatous lesion, rather than true neoplasm. Odontomas most of the times occur solitarily and at times are associated with impacted tooth and other odontogenic cysts and neoplasms, for which the identification and removal of the lesion is urged. [1,2,3]

Etiology of odontomas still remains unknown, however some studies have focused on the remnants of dental lamina as the major etiological factor. Some suggest that it is due to complete functional discrepancy of ameloblasts or odontoblasts occurs during the development of enamel and dentin, but in an irregular arrangement due to the lack of association of these cells to reach a usual state of morphodifferentiation. It might be congenital or due to an altered gene or postnatal genetic intervention of odontogenesis. [7] Genetic incongruities like Gardner’s syndrome, Hermann’s syndrome and local trauma, infectious or inflammatory processes, have been accredited to the development of odontomas. [1]

WHO classified odontoma into three variants namely, central, peripheral and erupted. Central or intraosseous variant is further subclassified into compound and complex, where compound odontoma presents as malformations in which dental tissues are well-organized in orderly pattern than in the complex odontoma, such that lesion consists of more teeth like structures. Complex odontoma is a malformation in which all dental tissues are well formed but occurring in less orderly pattern without anatomical resemblance to a tooth. [2,5]

Compound odontomas are painless benign lesion with limited growth potential compared to complex odontomas. It usually occur in children and young adult below the age range of 20 years with equal sex predilection. Maxillary anterior region is most frequently involved site. Radiographically, it is characterized by a radiopaque mass of different sizes,




Figure 1a: Radiograph showing radiopaque mass in 12 region



Figure 1b: Macroscopic feature showing small tooth like structure





Figure 2a: Radiopaque mass in the region of 11, obstructing the eruption pathway of 11. Impaction of 11 is also evident



Figure 2b: Macroscopic feature showing a hard tissue bits resembling teeth



Figure: 2c Photomicrograph of ground sectioned specimen showing fusion of teeth like structures containing well organized thin hypoplastic enamel, dentin cementum with incomplete root formation (4X)

Criteria	Case- 1	Case-2	Compound odontoma – Literature review
Age/ Sex	9/F	11/F	Childhood/adolescence (Below 20 years) M:F=1:1
Site	12	11	Maxillary anterior region (73%)
Clinical Presentation	Painless swelling with missing 12	Painless swelling with missing 11	Painless, non-aggressive lesion, often associated with unerupted /Impacted tooth
Radiological features	Radiopaque mass impacted 12	Radiopaque mass impacted 11	Radiopaque mass of multiple, small calcified structures with an anatomical similarity to normal teeth , usually surrounded by radiolucent zone
Microscopical features	Crown with well organized enamel and dentin like substances- No evidence of root	Thin, hypoplastic enamel, dentin & cementum -Fusion of 3 teeth like structures evident	Thin hypoplastic enamel, with mature tubular dentin and cementum resembling tooth structure
Treatment	Enucleation	Enucleation	Conservative surgical enucleation
Prognosis & follow up	No recurrence (5 month)	No recurrence (5 month)	Excellent prognosis No recurrence rate

Table 1: Showing comparison of features of 2 presented cases with the literature evidence

with variable amount of enamel, dentin, cementum and pulp and they resemble malformed or supernumerary teeth. The lesion is usually circumscribed by a narrow, radiolucent rim which corresponds to the fibrous capsule. [1,8,9]

Compound odontoma comprises of multiple structures such as enamel, dentin, cementum and even pulp chamber which resembles small teeth like structure, associated with this presence loose fibrous matrix were also present. Under decalcification, enamel that capped the teeth like structure were lost and leaves ample amount of enamel matrix. [5]

The conservative surgical excision of compound odontomas considered to be a treatment of choice. [8] Since odontomas are often associated with unerupted impacted teeth there could be a possibility of tooth eruption after surgical excision of the lesion. There is no recurrence reported till date, however the lesion may recur if it is incompletely removed at its early soft tissue stage.[3] Odontomas involving the maxillary antrum may lead to various complications such as orbital infections, epidural and subdural emphysema, meningitis, cavernous sinus thrombosis, brain abscess and death. [10]

This literature elaborates on two cases that reported to our OPD, with a complaint of missing upper anterior teeth. Both the cases upon examination and investigations were found to be compound odontoma. Our cases were in accordance with the literature with respect to age, radiographic findings, and clinical presentation. The macroscopic and microscopic features revealed structures resembling enamel, dentin and cementum. The lesions were enucleated conservatively and followed up. No recurrence was noticed during the follow up period. Earlier detection and treatment is emphasized to avoid complications that may arise in case of syndromic association and occurrence of an ameloblastic fibro odontoma.[11,12]

Longstanding compound odontoma associated with ameloblastic fibro odontoma [12] may further transform into ameloblastic fibro sarcoma which also occurs in the earlier period of life.[13]

References :

1. Bordini J Jr, Contar CM, Sarot JR, Fernandes A, Machado MA. Multiple compound odontomas in the jaw: case report

and analysis of the literature. *J Oral Maxillofac Surg.* 2008; 66(12):2617–20.

2. Rajendran R, Sivapathasundharam B. *Shafer’s Textbook of Oral Pathology: 7th ed.* India: Elsevier; 2012

3. Reichart PA, Philipsen HP. *Odontogenic tumours and Allied lesions.* London: Quintessence Publishing Co Ltd; 2004. pp. 150–53,

4. Cuesta SA , Albiol JG , Aytes LB, Escoda CG. Review of 61 cases of odontoma. Presentation of an erupted complex odontoma. *Med Oral.* 2003; 8(5):366-73

5. Neville BW, Damm DD, Allen CM, Bouquot J. *Oral and maxillofacial pathology.* 3rd ed. St. Louis, Missouri: Saunders Elsevier; 2009.

6. Gujjar PK , Sahni P, Pereira T, Zingade J. Multiple Compound Odontomas in the Jaws: A Rare Case Report. *Journal of Clinical and Diagnostic Research.* 2015;9(12): ZD05-ZD06

7. Serra-Serra G, Berini-Aytés L, Gay-Escoda C. Erupted odontomas: a report of three cases review of the literature. *Med Oral Patol Oral Cir Bucal.* 2009;14(6):E299–303.

8. Erdogan, Ztunc H, Evlice B, Altug HA, Günhan . Compound odontoma involving the four quadrants of the jaws: a case report and review of the literature. *Quintessence Int.* 2014; 45(4):341–44.

9. Ajike SO, Adekeye EO. Multiple compound odontomas in the facial bones: A case report. *Int J Oral Maxillofac Surg.* 2000;29 (6):443–44.

10. Guledgud MV, Degala S, Patil K, Keshari D. Multiple extensive complex odontomas of the jaws. *Int J Dent Sci Res.* 2014;2(6):128–32.

11. Hawkins PL, Sadeghi EM. Ameloblastic fibroodontoma: Report of case. *Journal of Oral and Maxillofacial Surgery.* 1986; 44: 1014-1019.

12. Slootweg PJ. An analysis of the interrelationship of the mixed odontogenic tumors-Ameloblastic fibroma, ameloblastic fibro-odontoma, and the odontomas. *Oral Surgery, Oral Medicine, Oral Pathology.* 1981; 51: 266-276.

13. Chang H, Shimizu MS, Precious DS. Ameloblastic Fibro-odontoma: A Case Report. *J Can Dent Assoc* 2002; 68(4):243-6

Address of Correspondence

Dr. Vidyalakshmi. S. M.D.S.,
Reader
Department of Oral Pathology and Microbiology
Indira Gandhi Institute of Dental Sciences
Sri Balaji Vidhyapeeth University
Pondicherry. India.
+91-9442503635
vidyan@igids.ac.in

Authors:

¹III year BDS, Indira Gandhi Institute of Dental Sciences, Sri Balaji Vidhyapeeth University, Pondicherry

²III year BDS, Indira Gandhi Institute of Dental Sciences, Sri Balaji Vidhyapeeth University, Pondicherry

³Post graduate student, Department of Oral Pathology and Microbiology, Indira Gandhi Institute of Dental Sciences, Sri Balaji Vidhyapeeth University, Pondicherry

⁴Reader, Department of Oral Pathology and Microbiology, Indira Gandhi Institute of Dental Sciences, Sri Balaji Vidhyapeeth University, Pondicherry.

How to cite this article :

Bharanippriyaa.V, Preethe.M, Umamaheswari.G, Vidyalakshmi.S. Odontoma- A Benign Mixed Odontogenic Tumor –Report of Two Cases. Journal of Scientific Dentistry, 2017;7(1):32-36

Source of Support : Nil, **Conflicts of Interest :** None declared